


SP-TOX

A companion of Toxic Management



Mycotoxins are secondary metabolites produced by certain moulds / fungi. Mycotoxicosis causes less overt signs but often is associated with hidden losses due to immunosuppression, hepatic & renal dysfunction, poor health & productivity. Toxin reduces the organs function such as liver & kidney. It also negatively impact on the poultry immune system & native defense mechanism.

SP TOX has an enormous surface area so it adsorbs wide range of toxins in the intestine. It is a multi-component toxin binder specially developed as a complete solution to the Mycotoxins problem. Selected ingredients binds wide range of various toxins presence in the Feed.

Key features:

- SP TOX formulated with selected high quality ingredients to perform as a broad Spectrum Mycotoxin absorbent in challenge condition.
- Its largest surface area to bind extreme and various toxins in wide range.
- The unique ingredients supports to control chemical toxicity due to pesticides and heavy metals.
- Selected silicates to adsorb both polar and dipolar Mycotoxins
- Maximizes Feed conversion and Improves overall performance
- SP TOX does not absorb vitamins, Minerals and other nutrients It reduces Mortality & Secondary bacterial diseases problem
- Protects the liver against the damage and mobilizes the fat accumulated in the liver

Benefits:

- To adsorb a wide range of mycotoxins.
- To protect and rehabilitates the liver and kidney.
- To reduce immunosuppression caused by mycotoxin.
- To Improve birds health through increased resistance to disease .
- To improve the immunity and feed intake.
- It formulated with selected high quantity ingredients to perform as a broad spectrum.
- It supports to control chemical toxicity due to pesticides and heavy metals.
- To maximize feed conversion and improves overall performance.

Composition:

Specially treat phyllosilicates, Highly observing tectosilicates, Mannan oligosaccharide (MOS) – Yeast cell wall, HSCAS, Activated carbon, Organic acids, Phytogetic derivative and blend of other toxin binding derivates.

Dosage :

Layer and Broiler : 0.5 Kg to 1 Kg per ton of feed at moisture level < 15%
1 Kg to 1.5 Kgs per ton of feed at moisture level > 15%

Cattle : 0.5 Kg to 1 Kg per ton of Feed.

Manufactured & Marketed by :



No. 1/258/3, Begapalli Vilage, Begapalli – Biderguppa Road, Hosur – 635126, Krishnagiri Distict, Tamilnadu, India.
Mobile : + 91 9790155126 E mail : gosanutribiotech@gmail.com Website : www.gosanutribiotech.com